

Stochastic background of relic scalar gravitational waves from scalar-tensor gravity

Salvatore Capozziello

Dipartimento di Scienze fisiche, Università di Napoli “ Federico II”, INFN Sezione di Napoli,
Compl. Univ. di Monte S. Angelo, Edificio G, Via Cinthia, I-80126, Napoli, Italy

Christian Corda

INFN Sezione di Pisa and Università di Pisa, Via F. Buonarroti 2, I-56127, Pisa,
Italy European Gravitational Observatory (EGO), Via E. Amaldi, I-56021 Pisa, Italy

Mariafelicia De Laurentis

Politecnico di Torino and INFN Sezione di Torino,
Corso Duca degli Abruzzi 24, I-10129 Torino, Italy

A stochastic background of relic gravitational waves is achieved by the so called adiabatically-amplified zero-point fluctuations process derived from early inflation. In principle, it provides a distinctive spectrum of relic gravitational waves. In the framework of scalar-tensor gravity, we discuss the scalar modes of gravitational waves and the primordial production of this scalar component which is generated beside tensorial one. We discuss also the upper limit for such a relic scalar component with respect to the WMAP constraints.

Keywords: Scalar-tensor gravity; gravitational waves; cosmology.

The design and the construction of a number of sensitive detectors for gravitational waves (GWs) is underway today. There are some laser interferometers like the VIRGO detector, built in Cascina, near Pisa, Italy, by a joint Italian-French collaboration, the GEO 600 detector built in Hanover, Germany, by a joint Anglo-German collaboration, the two LIGO detectors built in the United States (one in Hanford, Washington and the other in Livingston, Louisiana) by a joint Caltech-Mit collaboration, and the TAMA 300 detector, in Tokyo, Japan. Many bar detectors are currently in operation too, and several interferometers and bars are in a phase of planning and proposal stages (for the current status of gravitational waves experiments see [1, 2]). The results of these detectors will have a fundamental impact on astrophysics and gravitation physics. There will be a huge amount of experimental data to be analyzed, and theorists would face new aspects of physics from such a data stream. Furthermore, GW detectors will be of fundamental importance to probe the General Relativity or every alternative theory of gravitation [3, 4, 5, 6]. A possible target of these experiments is the so called stochastic background of gravitational waves [7, 8, 9, 10, 11]. The production of the primordial part of this stochastic background (relic GWs) is well known in literature starting from the works by [7] and [10], which, using the so called adiabatically-amplified zero-point fluctuations process, have shown in two different ways how the inflationary scenario for the early universe can, in principle, provide the signature for the spectrum of relic GWs.

In this letter, after a short summary of scalar-tensor gravity admitting the existence of GW scalar modes, the primordial production of this scalar component is discussed. Besides, the results are confronted to the WMAP data [12, 13] in order to achieve an upper limit for the scalar part of relic GWs.

Scalar-tensor gravity theories are a particular case of Extended Theories of Gravity which are revealing a useful paradigm to deal with several problems in cosmology, astrophysics and fundamental physics (for a comprehensive discussion see, for example [14, 15]). In the most general case, considering only the Ricci scalar among the curvature invariants, they arises from the action

$$S = \int d^4x \sqrt{-g} \left[F(R, \square R, \square^2 R, \square^k R, \phi) - \frac{\epsilon}{2} g^{\mu\nu} \phi_{;\mu} \phi_{;\nu} + \mathcal{L}_m \right], \quad (1)$$

where F is an unspecified function of curvature invariants and of a scalar field ϕ and $\square$ is the D'Alembert operator. The term $\mathcal{L}_m$ is the minimally coupled ordinary matter contribution. Scalar-tensor gravity, is recovered from (1) through the choice

$$F(R, \phi) = f(\phi)R - V(\phi), \quad \epsilon = -1. \quad (2)$$

Considering (2), a general action for scalar-tensor gravity in four dimensions is

$$S = \int d^4x \sqrt{-g} \left[f(\phi)R + \frac{1}{2} g^{\mu\nu} \phi_{;\mu} \phi_{;\nu} - V(\phi) + \mathcal{L}_m \right], \quad (3)$$

which can be recast in a Brans-Dicke-like form [16] by

$$\varphi = f(\phi), \quad \omega(\varphi) = \frac{f(\phi)}{2'f(\phi)}, \quad W(\varphi) = V(\phi(\varphi)) \quad (4)$$

and then

$$S = \int d^4x \sqrt{-g} \left[\varphi R - \frac{\omega(\varphi)}{\varphi} g^{\mu\nu} \varphi_{;\mu} \varphi_{;\nu} - W(\varphi) + \mathcal{L}_m \right]. \quad (5)$$

By varying the action (5) with respect to $g_{\mu\nu}$, we obtain the field equations

$$G_{\mu\nu} = -\frac{4\pi\tilde{G}}{\varphi} T_{\mu\nu}^{(m)} + \frac{\omega(\varphi)}{\varphi^2} (\varphi_{;\mu} \varphi_{;\nu} - \frac{1}{2} g_{\mu\nu} g^{\alpha\beta} \varphi_{;\alpha} \varphi_{;\beta}) + \frac{1}{\varphi} (\varphi_{;\mu\nu} - g_{\mu\nu} \square \varphi) + \frac{1}{2\varphi} g_{\mu\nu} W(\varphi) \quad (6)$$

while the variation with respect to φ gives the Klein - Gordon equation

$$\square \varphi = \frac{1}{2\omega(\varphi) + 3} \left[-4\pi\tilde{G} T^{(m)} + 2W(\varphi) + \varphi W'(\varphi) + \frac{d\omega(\varphi)}{d\varphi} g^{\mu\nu} \varphi_{;\mu} \varphi_{;\nu} \right]. \quad (7)$$

We are assuming physical units $G = 1$, $c = 1$ and $\hbar = 1$. $T_{\mu\nu}^{(m)}$ is the matter stress-energy tensor and $\tilde{G}$ is a dimensional, strictly positive, gravitational coupling constant [4, 5]. The Newton constant is replaced by the effective coupling

$$G_{eff} = -\frac{1}{2\varphi}, \quad (8)$$

which is, in general, different from G . General Relativity is recovered for

$$\varphi = \varphi_0 = -\frac{1}{2}. \quad (9)$$

In order to study gravitational waves, we assume first-order, small perturbations in vacuum ($T_{\mu\nu}^{(m)} = 0$). This means

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}, \quad \varphi = \varphi_0 + \delta\varphi \quad (10)$$

and

$$W \simeq \frac{1}{2} \alpha \delta\varphi^2 \Rightarrow W' \simeq \alpha \delta\varphi \quad (11)$$

for the self-interacting, scalar-field potential. These assumptions allow to derive the "linearized" curvature invariants $\tilde{R}_{\mu\nu\rho\sigma}$, $\tilde{R}_{\mu\nu}$ and $\tilde{R}$ which correspond to $R_{\mu\nu\rho\sigma}$, $R_{\mu\nu}$ and R , and then the linearized field equations [5, 17]

$$\begin{aligned} \tilde{R}_{\mu\nu} - \frac{\tilde{R}}{2} \eta_{\mu\nu} &= -\partial_\mu \partial_\nu \Phi + \eta_{\mu\nu} \square \Phi \\ \square \Phi &= m^2 \Phi, \end{aligned} \quad (12)$$

where

$$\Phi \equiv -\frac{\delta\varphi}{\varphi_0}, \quad m^2 \equiv \frac{\alpha\varphi_0}{2\omega + 3}. \quad (13)$$

The case $\omega = const$ and $W = 0$ has been analyzed in [5] considering the so-called "canonical" linearization [17]. In particular, the transverse-traceless (TT) gauge (see [17]) can be generalized to scalar-tensor gravity obtaining the total perturbation of a GW incoming in the $z+$ direction in this gauge as

$$h_{\mu\nu}(t-z) = A^+(t-z) e_{\mu\nu}^{(+)} + A^\times(t-z) e_{\mu\nu}^{(\times)} + \Phi(t-z) e_{\mu\nu}^{(s)}. \quad (14)$$

The term $A^+(t-z) e_{\mu\nu}^{(+)} + A^\times(t-z) e_{\mu\nu}^{(\times)}$ describes the two standard (i.e. tensorial) polarizations of a gravitational wave arising from General Relativity in the TT gauge [17], while the term $\Phi(t-z) e_{\mu\nu}^{(s)}$ is the extension of the TT

gauge to the scalar case. This means that, in scalar-tensor gravity, the scalar field generates a third component for the tensor polarization of GWs. This is because three different degrees of freedom are present (see Eq.(32) of [5]), while only two are present in standard General Relativity.

Let us now take into account the primordial physical process which gave rise to a characteristic spectrum Ω_{sgw} for the early stochastic background of relic scalar GWs. The production physical process has been analyzed, for example, in [7, 10, 18] but only for the first two tensorial components of eq. (14) due to standard General Relativity. Actually the process can be improved considering also the third scalar-tensor component.

Before starting with the analysis, it has to be emphasized that, considering a stochastic background of scalar GWs, it can be described in terms of the scalar field Φ and characterized by a dimensionless spectrum (see the analogous definition for tensorial waves in [7, 8, 9, 10])

$$\Omega_{sgw}(f) = \frac{1}{\rho_c} \frac{d\rho_{sgw}}{d \ln f}, \quad (15)$$

where

$$\rho_c \equiv \frac{3H_0^2}{8\pi G} \quad (16)$$

is the (actual) critical energy density of the Universe, H_0 the today observed Hubble expansion rate, and $d\rho_{sgw}$ is the energy density of the scalar part of the gravitational radiation contained in the frequency range f to $f + df$. We are considering now standard units.

The existence of a relic stochastic background of scalar GWs is a consequence of general assumptions. Essentially it derives from basic principles of Quantum Field Theory and General Relativity. The strong variations of gravitational field in the early Universe amplifies the zero-point quantum fluctuations and produces relic GWs. It is well known that the detection of relic GWs is the only way to learn about the evolution of the very early universe, up to the bounds of the Planck epoch and the initial singularity [7, 8, 9, 10, 11]. It is very important to stress the unavoidable and fundamental character of such a mechanism. It directly derives from the inflationary scenario [19, 20], which well fit the WMAP data in particular good agreement with almost exponential inflation and spectral index ≈ 1 , [12, 13].

A remarkable fact about the inflationary scenario is that it contains a natural mechanism which gives rise to perturbations for any field. It is important for our aims that such a mechanism provides also a distinctive spectrum for relic scalar GWs. These perturbations in inflationary cosmology arise from the most basic quantum mechanical effect: the uncertainty principle. In this way, the spectrum of relic GWs that we could detect today is nothing else but the adiabatically-amplified zero-point fluctuations [7, 10]. The calculation for a simple inflationary model can be performed for the scalar field component of eq. (14). Let us assume that the early Universe is described an inflationary de Sitter phase emerging in a radiation dominated phase [7, 8, 10]. The conformal metric element is

$$ds^2 = a^2(\eta)[-d\eta^2 + d\vec{x}^2 + h_{\mu\nu}(\eta, \vec{x})dx^\mu dx^\nu], \quad (17)$$

where, for a purely scalar GW the metric perturbation (14) reduces to

$$h_{\mu\nu} = \Phi e_{\mu\nu}^{(s)}, \quad (18)$$

Following [7, 10], in the de Sitter phase, we have:

$\eta < \eta_1$	conformal time
$P = -\rho$	equation of state
$\eta_1^2 \eta_0^{-1} (2\eta - \eta_0)^{-1}$	scale factor
$H_{ds} = c\eta_0/\eta_1^2$	Hubble constant

while, in the radiation dominated phase we have, respectively,

$\eta > \eta_1$	conformal time
$P = \rho/3$	equation of state
η/η_0	scale factor
$H = c\eta_0/\eta^2$	Hubble constant

η_1 is the inflation-radiation transition conformal time and η_0 is the value of conformal time today. If we express the scale factor in terms of comoving time $cdt = a(t)d\eta$, we have

$$a(t) \propto \exp(H_{ds}t), \quad a(t) \propto \sqrt{t} \quad (19)$$

for the de Sitter and radiation phases respectively. In order to solve the horizon and flatness problems, the condition $\frac{a(\eta_0)}{a(\eta_1)} > 10^{27}$ has to be satisfied. The relic scalar-tensor GWs are the weak perturbations $h_{\mu\nu}(\eta, \vec{x})$ of the metric (18) which can be written in the form

$$h_{\mu\nu} = e_{\mu\nu}^{(s)}(\hat{k})X(\eta)\exp(\vec{k} \cdot \vec{x}), \quad (20)$$

in terms of the conformal time η where $\vec{k}$ is a constant wavevector. From Eq.(20), the scalar component is

$$\Phi(\eta, \vec{k}, \vec{x}) = X(\eta)\exp(\vec{k} \cdot \vec{x}). \quad (21)$$

Assuming $Y(\eta) = a(\eta)X(\eta)$, from the Klein-Gordon equation in the FRW metric, one gets

$$Y'' + (|\vec{k}|^2 - \frac{a''}{a})Y = 0 \quad (22)$$

where the prime ' denotes derivative with respect to the conformal time. The solutions of Eq. (22) can be expressed in terms of Hankel functions in both the inflationary and radiation dominated eras, that is:
For $\eta < \eta_1$

$$X(\eta) = \frac{a(\eta_1)}{a(\eta)}[1 + H_{ds}\omega^{-1}]\exp -ik(\eta - \eta_1), \quad (23)$$

for $\eta > \eta_1$

$$X(\eta) = \frac{a(\eta_1)}{a(\eta)}[\alpha \exp -ik(\eta - \eta_1) + \beta \exp ik(\eta - \eta_1)], \quad (24)$$

where $\omega = ck/a$ is the angular frequency of the wave (which is function of the time being $k = |\vec{k}|$ constant), α and β are time-independent constants which we can obtain demanding that both X and $dX/d\eta$ are continuous at the boundary $\eta = \eta_1$ between the inflationary and the radiation dominated eras. By this constraint, we obtain

$$\alpha = 1 + i\frac{\sqrt{H_{ds}H_0}}{\omega} - \frac{H_{ds}H_0}{2\omega^2}, \quad \beta = \frac{H_{ds}H_0}{2\omega^2} \quad (25)$$

In Eqs. (25), $\omega = ck/a(\eta_0)$ is the angular frequency as observed today, $H_0 = c/\eta_0$ is the Hubble expansion rate as observed today. Such calculations are referred in literature as the Bogoliubov coefficient methods [7, 10].

In an inflationary scenario, every classical or macroscopic perturbation is damped out by the inflation, i.e. the minimum allowed level of fluctuations is that required by the uncertainty principle. The solution (23) corresponds to a de Sitter vacuum state. If the period of inflation is long enough, the today observable properties of the Universe should be indistinguishable from the properties of a Universe started in the de Sitter vacuum state. In the radiation dominated phase, the eigenmodes which describe particles are the coefficients of α , and these which describe antiparticles are the coefficients of β (see also [21]). Thus, the number of particles created at angular frequency ω in the radiation dominated phase is given by

$$N_\omega = |\beta_\omega|^2 = \left(\frac{H_{ds}H_0}{2\omega^2}\right)^2. \quad (26)$$

Now it is possible to write an expression for the energy density of the stochastic scalar relic gravitons background in the frequency interval $(\omega, \omega + d\omega)$ as

$$d\rho_{sgw} = 2\hbar\omega \left(\frac{\omega^2 d\omega}{2\pi^2 c^3}\right) N_\omega = \frac{\hbar H_{ds}^2 H_0^2}{4\pi^2 c^3} \frac{d\omega}{\omega} = \frac{\hbar H_{ds}^2 H_0^2}{4\pi^2 c^3} \frac{df}{f}, \quad (27)$$




Figure 1: The spectrum of relic scalar GWs in inflationary models is flat over a wide range of frequencies. The horizontal axis is $\log_{10}$ of frequency, in Hz. The vertical axis is $\log_{10} \Omega_{sgw}$. The inflationary spectrum rises quickly at low frequencies (wave which re-entered in the Hubble sphere after the Universe became matter dominated) and falls off above the (appropriately redshifted) frequency scale f_{max} associated with the fastest characteristic time of the phase transition at the end of inflation. The amplitude of the flat region depends only on the energy density during the inflationary stage; we have chosen the largest amplitude consistent with the WMAP constraints on scalar perturbations. This means that at LIGO and LISA frequencies, $\Omega_{sgw} < 2.3 * 10^{-12}$

where f , as above, is the frequency in standard comoving time. Eq. (27) can be rewritten in terms of the today and de Sitter value of energy density being

$$H_0 = \frac{8\pi G \rho_c}{3c^2}, \quad H_{ds} = \frac{8\pi G \rho_{ds}}{3c^2}. \quad (28)$$

Introducing the Planck density $\rho_{Planck} = \frac{c^7}{\hbar G^2}$ the spectrum is given by

$$\Omega_{sgw}(f) = \frac{1}{\rho_c} \frac{d\rho_{sgw}}{d \ln f} = \frac{f}{\rho_c} \frac{d\rho_{sgw}}{df} = \frac{16}{9} \frac{\rho_{ds}}{\rho_{Planck}}. \quad (29)$$

At this point, some comments are in order. First of all, such a calculation works for a simplified model that does not include the matter dominated era. If also such an era is also included, the redshift at equivalence epoch has to be considered. Taking into account also results in [18], we get

$$\Omega_{sgw}(f) = \frac{16}{9} \frac{\rho_{ds}}{\rho_{Planck}} (1 + z_{eq})^{-1}, \quad (30)$$

for the waves which, at the epoch in which the Universe becomes matter dominated, have a frequency higher than H_{eq} , the Hubble parameter at equivalence. This situation corresponds to frequencies $f > (1 + z_{eq})^{1/2} H_0$. The redshift correction in Eq.(30) is needed since the today observed Hubble parameter H_0 would result different without a matter dominated contribution. At lower frequencies, the spectrum is given by [7, 10]

$$\Omega_{sgw}(f) \propto f^{-2}. \quad (31)$$

As a further consideration, let us note that the results (29) and (30), which are not frequency dependent, does not work correctly in all the range of physical frequencies. For waves with frequencies less than today observed H_0 , the

notion of energy density has no sense, since the wavelength becomes longer than the Hubble scale of the Universe. In analogous way, at high frequencies, there is a maximal frequency above which the spectrum rapidly drops to zero. In the above calculation, the simple assumption that the phase transition from the inflationary to the radiation dominated epoch is instantaneous has been made. In the physical Universe, this process occurs over some time scale $\Delta\tau$, being

$$f_{max} = \frac{a(t_1)}{a(t_0)} \frac{1}{\Delta\tau}, \quad (32)$$

which is the redshifted rate of the transition. In any case, Ω_{sgw} drops rapidly. The two cutoffs at low and high frequencies for the spectrum guarantee that the total energy density of the relic scalar gravitons is finite. For GUT energy-scale inflation it is of the order [7]

$$\frac{\rho_{ds}}{\rho_{Planck}} \approx 10^{-12}. \quad (33)$$

These results can be quantitatively constrained considering the recent WMAP release. In fact, it is well known that WMAP observations put strongly severe restrictions on the spectrum. In Fig. 1 the spectrum Ω_{sgw} is mapped : considering the ratio ρ_{ds}/ρ_{Planck} , the relic scalar GW spectrum seems consistent with the WMAP constraints on scalar perturbations. Nevertheless, since the spectrum falls off $\propto f^{-2}$ at low frequencies, this means that today, at LIGO-VIRGO and LISA frequencies (indicated in fig. 1), one gets

$$\Omega_{sgw}(f)h_{100}^2 < 2.3 \times 10^{-12}. \quad (34)$$

It is interesting to calculate the corresponding strain at $\approx 100Hz$, where interferometers like VIRGO and LIGO reach a maximum in sensitivity. The well known equation for the characteristic amplitude [7, 10, 11] adapted to the scalar component of GWs can be used:

$$\Phi_c(f) \simeq 1.26 \times 10^{-18} \left(\frac{1Hz}{f}\right) \sqrt{h_{100}^2 \Omega_{sgw}(f)}, \quad (35)$$

and then we obtain

$$\Phi_c(100Hz) < 2 \times 10^{-26}. \quad (36)$$

Then, since we expect a sensitivity of the order of 10^{-22} for the above interferometers at $\approx 100Hz$, we need to gain four order of magnitude. Let us analyze the situation also at smaller frequencies. The sensitivity of the VIRGO interferometer is of the order of 10^{-21} at $\approx 10Hz$ and in that case it is

$$\Phi_c(100Hz) < 2 \times 10^{-25}. \quad (37)$$

The sensitivity of the LISA interferometer will be of the order of 10^{-22} at $\approx 10^{-3}Hz$ and in that case it is

$$\Phi_c(100Hz) < 2 \times 10^{-21}. \quad (38)$$

This means that a stochastic background of relic scalar GWs could be, in principle, detected by the LISA interferometer.

In summary, the above results point out that a further scalar component of GWs, coming from Extended Theories of Gravity, should be seriously considered in the signal detection of interferometers. As already discussed in [21], this fact could constitute an independent test for alternative theories of gravity or a further probe for General Relativity capable of ruling out other theories.

-
- [1] http://www.ligo.org/pdf_public/camp.pdf.
 - [2] http://www.ligo.org/pdf_public/hough02.pdf.
 - [3] S. Capozziello and A. Troisi, Phys. Rev. D 72, 044022 (2005).
 - [4] S. Capozziello - *Newtonian Limit of Extended Theories of Gravity in Quantum Gravity Research Trends* Ed. A. Reimer, pp. 227-276 Nova Science Publishers Inc., NY (2005) arXiv:gr-qc/0412088.
 - [5] S. Capozziello and C. Corda - Int. J. Mod. Phys. D 15, 1119 -1150 (2006).
 - [6] M. E. Tobar , T. Suzuki and K. Kuroda Phys. Rev. D 59, 102002 (1999).

- [7] B. Allen -Proceedings of the Les Houches School on Astrophysical Sources of Gravitational Waves, eds. Jean-Alain Marck and Jean-Pierre Lasota (Cambridge University Press, Cambridge, England 1998).
- [8] B. Allen and A.C. Ottewill - Phys. Rev. D 56, 545-563 (1997).
- [9] M. Maggiore - Phys. Rep. 331, 283-367 (2000).
- [10] L. Grishchuk et al. - Phys. Usp. 44 1-51 (2001); Usp.Fiz.Nauk 171 3-59 (2001).
- [11] C. Corda - *VIRGO Report: "Relic gravitational waves: a "snapshot" of the primordial universe"* VIRGO-NOTE-PIS 1390-237 (2003) - www.virgo.infn.it/Documents.
- [12] C.L. Bennet et al. - ApJS 148, 1.
- [13] D.N. Spergel et al. - ApJS 148, 195.
- [14] S. Nojiri and S.D. Odintsov *Int. J. Geom. Meth. Mod. Phys.* **4**, 115 (2007), arXiv: hep-th/0601213.
- [15] S. Capozziello and M. Francaviglia, arXiv: 0706.1146 [astro-ph] (2007).
- [16] C. Brans and R. H. Dicke - Phys. Rev. 124, 925 (1961).
- [17] C. W. Misner , K. S. Thorne and J. A. Wheeler - "Gravitation" - W. H. Feeman and Company 1973.
- [18] B. Allen - Phys. Rev. D 3-7,2078 (1988).
- [19] G.S. Watson- "An exposition on inflationary cosmology" - North Carolina University Press (2000).
- [20] A. Guth - Phys. Rev. 23, 347 (1981).
- [21] S. Capozziello, Ch. Corda, M. De Laurentis, Mod. Phys. Lett. **A 22**, 1097 (2007).